

Bangkok Ending AIDS 2073

Reduced new HIV infection

Reduced deaths related to AIDS

Reduced discrimination

Reach and Recruit

1. Reach and recruit 90% of key populations for comprehensive prevention services

Provide knowledge and create awareness

- *Peer
- *outreach
- *Drop-in center
- *Pharmacy networks
- *Social Media
- *Public media
- *STI screening and treatment

Equipment

Sufficient condoms and lubricants provided in the field

Test

2. Promote HCT and combination prevention of HIV with the focus to cover 80 % of key populations

Counselling and Testing

- *Proactive service (Mobile VCT)
- *HIV testing with same day result
- * VCT/PICT
- *Referral system

Treat

3. Provide ARV treatment to 85% of all persons living with HIV

Develop friendly-service system

- *Care and treatment management
- *CD4 count and viral load testing service
- *Support referral systems and networking
- *Expand laboratory networks (CD4, viral load, Hepatitis C)
- *TB & HIV treatment service
- *Pre-ARV Treatment Counseling
- *Access to service systems and support standards

Retain

4. No less than 90% continue in the service system and adhere to ARV

Social media and measures

- *Work through community leaders to retain both negative and positive persons in the prevention and treatment system
- *Management by the community
- *Strengthen care through peer helpers

Reduce stigma, discrimination and create a legal environment that supports the implementation of the strategy.

Management: Provide technical support, create and utilize strategic information.

Target groups include youth, MSM, sex workers, persons who inject drugs and migrants.


Civil Society

Private Sector

Government Sector beyond BMA

Government Sector governed by BMA

Involvement of all partner networks and sectors.



FAST-TRACK

ENDING THE AIDS EPIDEMIC BY 2030

90-81-90 global indicator definitions



diagnosed

Indicator 1

90% diagnosed
of all people living with HIV



on treatment

Indicator 2

81%
90% on treatment
~~of all people diagnosed~~
of all people living with HIV




virally suppressed

Indicator 3

90% virally suppressed
of all people on treatment

GARPR 2016 guidelines and indicator descriptions will reflect modifications to 90-90-90 target as described in WHO's Consolidated SI Guidelines

Mapping Hot Spot



Legend
Number of target in districts

Number

- ยังไม่ได้ทำ Mapping
- 1-500 คน
- 501-1000 คน
- 1001-1500 คน
- 1501-2000 คน
- 2001-2500 คน
- 2501-3000 คน
- มากกว่า 3000 คน



Integrated Behavioral Surveillance Survey among MSM in Bangkok


Objective

- 1. To determine the prevalence of infection with HIV and the behaviors associated with HIV infection among men who have sex with men.**
- 2. Epidemiological trends of prevalence of HIV infection and the behaviors associated with HIV infection among men who have sex with men.**
- 3. Application procedures for monitoring behavior and prevalence of HIV infection among men who have sex with men to perform the normal system.**

Study Designs

Meeting Planners

Place



Ratchathewi

Bangkapi

Wattana

Wang Thong Lang

Phayathai

Chatuchak

Phra Nakhon

Huai Khwang

Park

Pub

Sauna

Study Designs

Time

April –
May 2016



Person

MSM 297

MSW 150

TG 150




Material

- Consent Form
- Oraquick
- Questionnaire by smart phone

Methodology

Contract
NGO
actions







Result & Discussion

Demographics Data



Age

< 25 year




> 25 year




Demographics Data



Result Prevalence 2016





HIV Knowledge (UNGASS)




1. If the partners are the only ones who are not infected with HIV can reduced the risk of infection with HIV / AIDS.
2. Use a condom every time during sexual intercourse can reduce the risk of infection by HIV / AIDS.
3. People who look healthy are infected with HIV / AIDS.
4. People can be infected with HIV / AIDS from eating infected with HIV / AIDS.
5. People can be infected with HIV / AIDS from mosquito bites.


Use of condoms among male-male sex with male regular partner



Source : Integrated Biological Behavioral Surveillance (IBBS)




Access to prevention information/supplies among MSM




Source : Integrated Biological Behavioral Surveillance (IBBS)

Access to prevention information/supplies among TG



Source : Integrated Biological Behavioral Surveillance (IBBS)

Access to prevention information/supplies among MSM



SOURCE : มูลนิธิรณรงค์ฯไทย, RSAT , SWING, บางกอกเรนโบว์, NGOกทม. , หน่วยงานสังกัดกรุงเทพมหานคร



 **Conclusion**


Conclusion

1. **New infection is more among MSM**
2. **MSM mainly younger**
3. **MSM mainly an educational degree**
4. **Prevalance of infection among MSM declined from 2014**
5. **Bar have prevalance of infection among MSM in 2016**
6. **Condom use and Access to prevention information/supplies among MSM and TG under MSW**



Bangkok Action Plan

Bangkok Fast Track towards 90-90-90 targets for MSM/TGW and MSW response

1. Access to prevention among MSM at high risk and younger
2. Development of network synchronization in all sector
3. Increasing access by MSM and TG women to HCT,STI screening and PrEP in BMA health center
4. Prototy  HIV and testing are normalise

